

## CONTENTS

## ARTICLES

IR and <sup>1</sup> H NMR Spectra of Tin(II) Complexes with 1-Substituted Tetrazoline-5-thiones R.N. PANDEY	1657
Low-valent Organometallic Derivatives of Palladium(O), Platinum(O) and Rhodium(I) with 4-Amino-5-mercapto-3-substituted-1,2,4-triazoles R.N. PANDEY, ASHOK KUMAR and D.P. SINGH	1661
Biological Excess Sludge Reduction in Municipal Wastewater Treatment by Chlorine AFSHIN TAKDASTAN, ALI AKBAR AZIMI and NEMATOLLAH JAAFARZADEH	1665
Carbon Dioxide Reduction Options in Power Generation SHIV PRATAP RAGHUVANSHI, RENU SINGH, A.K. RAGHAV and A. CHANDRA	1675
Antioxidant Activity and Brine Shrimp Lethality of <i>Tragia involucrata</i> L. MANTENA G. SAVITHRI, BHUPATIRAJU A.S. CHANDANADEV, ALLURI V. KRISHNARAJU, CHIRRAVURI V. RAO and GOLAKOTI TRIMURTULU	1684
Synthesis and Crystal Structure of 1-(2-(1 <i>H</i> -Benzoimidazol-1-yl)ethyl)-1 <i>H</i> -benzoimidazol-3-ium Chloride Hydrate JUN YANG, CAI FENG BI, TINGTING YOU and YUHUA FAN	1689
Effect of Agitation on the Extraction of Palladium from Spent Industrial Catalyst with Supercritical Carbon Dioxide MUHAMMAD FAISAL, YOICHI ATSUTA and HIROYUKI DAIMON	1693
Biosorption of Lead and Nickel by <i>Medicago sativa</i> (alfalfa) and <i>Datura</i> from Contaminated Solution S.B. MOSAVI, S. KARIMI and V. FEIZIASL	1700
A New Lignan from the Fruit of <i>Schisandra propinqua</i> var. <i>propinqua</i> and its Anti-HIV Activity PENG FAN, XIN ZHANG, TAO LI, YUN DAI, GAN-PENG LI and QIU-FEN HU	1705
Interaction Studies of <i>Bis</i> -( <i>p</i> -aminobenzoato)diaquanickel(II) and <i>Bis</i> -( <i>p</i> -aminobenzoato)triaquacadmium(II) Complexes with Cat Genomic DNA ALI ARSLANTAS, MAHMUT SOZMEN and HACALI NECEFOGLU	1709
Spectrophotometric Determination of Cu(II) using 2-Hydroxy-4 <i>N</i> -butoxy-5-bromo Acetophenone Oxime S.S. SHAH and R.G. PARMAR	1717
Preparation and Evaluation of Starch Acetate Coated Microcapsules of Glipizide for Controlled Release K.P.R. CHOWDARY, P. SEENIVASAN and C. UMAMAHESWARA REDDY	1723
Determination of Vanadium(V) by Kinetic-Catalytic Spectrophotometric Method Using the Oxidation of Commassive Violet R 150 by Bromate MOHSEN KEYVANFARD	1729
Determination of 2,3,5-Triphenyltetrazolium Chloride Based on Flavinic Acid and 5-Nitrobarbituric acid Ion Pair Membrane A.G. FATHI EL SAID, N.M.H. RIZK and A.B. HAMDY MATTER	1736

Transition Metal Complexation Studies on Heterocyclic Ligands R.T. VASHI and C.D. SHELAT	1745
Investigation of Minimization of Excess Sludge Production in Sequencing Batch Reactor by Heating Some Sludge MARYAM PAZOKI, AFSHIN TAKDASTAN and NEMATOLLAH JAAFARZADEH	1751
Synthesis and Characterization of Transition Metal Complexes with 7-(R)-2-Amino-2-(4-hydroxyphenyl)acetamido]-3-methyl-3-cephem-4-carboxylic Acid Monohydrate J.H. DESHMUKH and M.N. DESHPANDE	1760
Spectrophotometric Determination of Moprolol in Pure Form and Pharmaceutical Formulations REKHA RAJEEV and RAJEEV KUMAR	1769
Synthesis and Characterization of Some Novel Complexes of Iron(III) with Ketoanil Schiff Base PRINCE PATHAK, MONAL SINGH and VISHAL PATHAK	1773
Composition of Leaf Essential Oil of <i>Eucalyptus camaldulensis</i> MUHAMMAD ASHRAF, QASIM ALI, FAROOQ ANWAR and ABDULLAH IJAZ HUSSAIN	1779
Effect of Solvent on Kinetics of Reduction of Methyl Violet with Potassium Iodide at Different Temperatures R. SAEED, M. ASHFAQ and S. RAZI KHAN	1787
Ethanolamines as Corrosion Inhibitors for Zinc in Sulfamic Acid V.A. CHAMPANERI and R.T. VASHI	1799
Extraction and Composition of Volatiles from Concrete of <i>Pandanus fascicularis</i> Lam PRASANT K. ROUT, SATYANARAYAN NAIK and Y. RAMACHANDRA RAO	1808
Synthesis, Characterization and Thermal Studies of a New Imine-oxime Ligand and its Cu <sup>2+</sup> , Ni <sup>2+</sup> , Co <sup>3+</sup> and Zn <sup>2+</sup> Complexes YUNUS KAYA, GAZI IREZ and HASENE MUTLU	1813
FT-IR, TG, XRD and SEM Studies of Activated Carbons Prepared From Agricultural Waste B.V. RAJESWARA REDDY, CH. CHAKRAPANI, CH. SURESH BABU, K. SOMASEKHARA RAO, KAVITA AGARWAL and MAHENDER PRASAD	1822
Determination of Molybdenum in Soil, Coal and Industrial Residue Samples by Flame Atomic Absorption Spectrometry ERGUL YIGMATEPE, HALIM AVCI and MEHMET YAMAN	1829
New Strategy for the Synthesis of (±) Cherylline Dimethyl Ether S. VENUGOPAL, J. RAMANATHAM, N. DEVANNA, A. SANJEEV KUMAR, SAMIR GHOSH, R. SOUNDARARAJAN, BHIMA KALE and G.N. MEHTA	1835
Improved Synthesis of Novel 2,4-Diamino-5-furfurylpyrimidine in Presence of Molecular Sieves R. RAJAPANDI, JAGADISH SINGH and T.K. MAITY	1841
Chemical Composition and Fatty Acid Contents of Chestnuts Grown in Bursa Province, Turkey SERKAN ÖZTÜRK, MURAT ÇETIN and MEHMET ÇETIN	1845
Synthesis of Tricyclic Derivatives of Antitumor Alkaloid Ellipticine YAVUZ ERGÜN, SIBEL GÜLLE, ADIL ÇAGDAS BILICI, MUSTAFA GÖÇMENTÜRK and GÜROL OKAY	1853

Evaluation of Chlorogenic Acid in <i>Flos lonicerae</i> by HPLC Method LIAO WEIPING and RUI YUKUI	1859
Thermodynamic Deprotonation of N,N'-Disubstituted Violuric Acid and Thiovioluric Acid in Dioxane-Water Mixtures DEEP SORABH SINGH and B.R. SINGH	1864
Synthesis and Characterization of Phenolic Resole Resins for Adhesive Applications FARIDEH GOURANLOU	1873
Occurrence of Aflatoxin M <sub>1</sub> in Raw Milk in Trakya Region, Turkey A. OZSUNAR, T. GUMUS, M. ARICI and M. DEMIRCI	1879
Estimation of Arsenic(III) by Adsorption and Filtration from Wastewater of Kanti Thermal Power, Muzaffarpur (India) KALPNA KUMARI	1885
Macroelement and Microelement Contents in Wild Edible Mushroom Species from the Southern Regions of China X.H. CHEN, H.B. ZHOU and G.Z. QIU	1889
Formulation and Statistical Optimization of Controlled Release Matrix Tablet of Zidovudine R.K. KAR, S. MOHAPATRA, P. MOHAPATRA, S.B. BHANJA and B.B. BARIK	1896
Exo-axial, Endo-axial and Exo-equatorial Anomeric Effects for Mono and Dihalo[1,4]dioxane in Solvent and Gas Phases S. ARSHADI, Y. SHAD, J. EMAMI, A.R. BEKHRADNIA, O. YAZDANI, R. PAHLEVAN, F. QAJAR, M. RAJAEI, A. BADELI and M.A. EBRAHIMZADEH	1907
Kinetics and Mechanism of Oxidation of $\beta$ -Benzoyl Propionic Acids by Pyridinium Chlorochromate in Aqueous Acetic Acid Medium P. SAHAYA AMUTHA and N. MATHIYALAGAN	1918
Fischer-Tropsch Synthesis on Co/SiO <sub>2</sub> Catalysts with Potassium Promoter HOSSEIN ATASHI, MEHDI SAHEBI-SHAHEMABADI and REZA AKHAVAN	1925
Thermal Properties and Gas Decomposition Products of Hafnium(IV) Acetylacetonate QIAOWEN CHANG, NI ZHANG, WEIPING LIU, QINGSONG YE, YAO YU, XIZHU CHEN and CHUNYING LI	1935
Effect of Saffron Extract on Proteins Biochemical Parameter of Serum MEHRDAD MODARESI, MANOCHEHR MESSRIPOUR and MEHRAN ASADI MARGHMALEKI	1939
Optical and Printability Properties of 2-[[4-(Aminosulfonyl)-phenyl]hydrazono]-N-(4-ethoxyphenyl)-3-oxobutanamide MEHMET OKTAV, SEVIM ROLLAS and EFE N. GENCOGLU	1944
Synthesis and Multinuclear NMR Study of Novel Complexes of Zn(II) and Hg(II) Containing Phosphorus Ylides SEYYED JAVAD SABOUNCHEI, MOHSEN AHMADI GHARACHEH and MARJAN HOSSEINZADEH	1949
Binding Constant: Fluorescence Quenching of Ciprofloxacin with Fe(III) and Zn(II) K.S. SIDDIQI, AYAZ MOHD, AFTAB ASLAM, PARWAZ KHAN and SHAISTA BANO	1957
Full Non-Rigid Group Theory for Hexamethylethane MOHAMMAD REZA DARAFSHEH and ALI MOGHANI	1966
Synthesis of Aryl-bis(3-methyl-1-phenyl-5-pyrazolone-4-yl)methane in Water AI-QING WANG, TONG-SHOU JIN, ZHAO-LI CHENG and TONG-SHUANG LI	1973

A Clean and Efficient Method for the Synthesis of 3-Aroyl-2,4,6-triaryl-4-hydroxy-1,1-cyclohexanedicarbonitriles in Water	1977
AI-QING WANG, TONG-SHOU JIN, LI-BIN LIU, ZHAO-LI CHENG and TONG-SHUANG LI	
Solubility of Edible Films from Amaranth Flour Enzymatic Hydrolyzate Modified with Dialdehyde Starch	1982
P. MOKREJS, J. HRNCIRIK, D. JANACOVA, V. VASEK and R. CERMAK	
Comparative Studies on Adsorption of Congo Red (CR) by Low Cost Adsorbents Prepared from Different Varieties of Melon Seeds	1993
NIDA IQBAL, MUHAMMAD IMRAN, JAVED IQBAL and ZAID MAHMOOD	
Adsorption Behaviour of XDA-200 Resin Toward <i>p</i> -Nitrophenol in Aqueous Solution	2003
ZHANWANG ZHENG, CHUNHUA XIONG and CAIPING YAO	
Catalase and Superoxide Dismutase Enzymes Activities in Flaxseed ( <i>Linum Usitatissimum</i> ) Extract	2013
GUZIDE YUCEBILGIC and YURDANUR BOZDEMIR	
Solid Phase Extraction and Graphite furnace Atomic Absorption Spectrometry for the Determination of Palladium	2018
ZHANGJIE HUANG, JUN CHANG, SHAOPING FENG and ZHONGQI YANG	
Effect of pH Variation on Solubility of Seed Protein of <i>Pisum sativum</i>	2027
G. NAGA BHUSHANA RAO and S.K. SHRIVASTAVA	
Synthesis and Characterization of Some Transition Metal Complexes of Dithiocarbamate Ligand Derived from <i>p</i> -Toluidine	2031
MOHAMMAD TARIQUE and MOHAMMAD ASLAM	
Stability Constants of Ternary Complexes of Cd(II) with Itaconic Acid and Some Amino Acids: A Polarographic Approach	2035
CHANCHAL KARADIA and O.D. GUPTA	
Electrochemical Studies of Copper(II) Complexes with Oxoglutarate at Dropping Mercury Electrode in Aqueous Medium	2043
O.D. GUPTA, MEENA and JYOTI SINGH	
Stability-Indicating HPLC Method for the Determination of Oxcarbazepine in Pharmaceutical Formulation	2051
U. BHAUMIK, A. BOSE, B. CHATTERJEE, A. GHOSH, P. SENGUPTA, S. AGARWAL, A. DAS and T.K. PAL	
Michael Adducts of 2-Methoxycarbonyl-1,4-benzoquinone with Different Donor Molecules	2058
A. ASHNAGAR and J.M. BRUCE	
A Suggested Improvement to The Plant Recovery Formula: A Case Study for Elazig Ferrochrome Plant	2073
HUSEYIN ANKARA	
Simultaneous Removal of Indomethacine, Papaverine and Allopurinol from Aqueous Solution by Using Submerged Aquatic Plant <i>Nasturtium officinale</i>	2081
VEYSI OKUMUS, ELIF ORAL, DAVUT BASARAN and AHMET ONAY	
Kinetics and Mechanism of Oxidation of <i>m</i> -Anisidine by Periodate Ion	2090
R.D. KAUSHIK, RAJDEEP MALIK, MANILA and AJAY KUMAR	

Synthesis and Pharmacological Screening of Indan-1-propionic Acids as Anti-hypercholesterolemic Agents M.R. ADAK	2099
Development of a Monitoring Method for Oxidative Coupling Reaction of 2-Naphthol in Solid State KEIVAN SHAYESTEH, JAFARSADEGH MOGHADDAS, MOHAMMAD HAGHIGHI and HABIBOLLAH ESKANDARI	2106
Chemical Synthesis of UDP-Glc <sub>4</sub> A from N-Acetyl Chondrosine and UDP-[β-D-Gal-(1→4)-α-D-Glu] from Lactose HAMMED H.A.M. HASSAN	2117
Polarographic Determination of Topiramate in Some Pharmaceuticals H. MANDIL, A.A. SAKUR and S. ALULU	2129
Kinetic-Spectrophotometric Studies on Periodate Oxidation of 2-Amino- <i>p</i> -Xylene in Acetone-Water Medium R.D. KAUSHIK, PURNIMA SUNDRIYAL, TARUN KUMAR and PRABHA SINGH	2136
Methyl Cellulose Based Sustained Release Thermosensitive <i>in situ</i> Fast Gelling Ocular Delivery of Ketorolac Tromethamine MANAS BHOWMIK, SANCHITA DAS, JAYDEEP SINHA, SWARNENDU BAG, DIPANKAR CHATTOPADHYAY and LAKSHMI K. GHOSH	2147
Spectrophotometric Study of Kinetics and Mechanism of Oxidation of <i>o</i> -Ethoxyaniline by Periodate Ion R.D. KAUSHIK, DHARMENDRA KUMAR, AJAY KUMAR and RAJDEEP MALIK	2155
Congo Red as a Modifier for Amperometric Determination of Ascorbic Acid R.E. SABZI, E. MINAEI and K. FARHADI	2165
Solid Phase Extraction and Spectrophotometric Method for Determination of Trace Gold with Octyl Benzimidazole Sulfide ZHANGJIE HUANG, JUN ZHANG, JUN CHANG, SHAOPING FENG and ZHONGQI YANG	2172
Synthesis and Characterization of Gold(III) Complexes with <i>bis</i> -(1,4-Sodium thioglycolate)butane Ligand EID A. ABDALRAZAQ, NABEEL H. BUTTRUS and ABD AL-MOUJOD M. ABD AL-RAHMAN	2179
Determination of Aluminum, Copper and Lead in Aleppo Company of Cable Industry (Syria) Using Industrial TSP, PM <sub>10</sub> and Chemical Trap Methods ABDUL AZIZ RAMADAN, HASNA MANDIL and MAHMOUD HASAN ISMAIL	2187
Determination of Arsenic and Some Heavy Metals Levels in Roadside Soils and Leaves of Aleppo City, Syria ABDUL AZIZ RAMADAN and HASNA MANDIL	2195
Determination of Epoxy Groups on γ-Glycidoxypolytrimethoxy-silane Doped Nanosilicas by Oscillo-polarographic DE-LIANG LI, LING ZHANG, ZHI-XIAN CHANG and ZHI-JUN ZHANG	2207
Effect of Mordant Methods on Silk Yarn: Dyeing with Natural Red Dyes and Improvement of Adsorbed Dyeability HOSSEIN NAJAFI and REZA ASSEFIPOUR	2214

Biotechnological Processes for The Elimination of Hydrogen Peroxide After Bleaching of Cotton/Jute Fabrics	
HOSSEIN NAJAFI and REZA ASSEFIPOUR	2221
Modeling the Tensile Properties of Soybean Protein Yarns	
HOSSEIN NAJAFI and REZA ASSEFIPOUR	2225
Preparation of Carbon Nanotubes/TiO <sub>2</sub> Composites with Multi-Walled Carbon Nanotubes and Titanium Alkoxides by Solvent Effect and Their Photocatalytic Activity	
W.C. OH, F.J. ZHANG and M.L. CHEN	2231
Studies on Corrosion of Metals in Surat Urban Environment	
R.T. VASHI, R.N. PATEL, K.S. CONTRACTOR and S.A. DESAI	2244
Geochemical Studies on the Role of Diadochy in the Cations Distribution in Minerals of Amphibole Group in Earth's Crust	
H.C. RAI, MEERA SHARMA, ANNAPURNA SHARMA, SARITA KUMARI, REETA KUMARI and ANITA KUMARI	2251
Synthesis, Spectroscopic and Electrochemical Studies of Cobalt(II) and Nickel(II) Macrocylic Complexes Derived from Oxalodihydrazide and Aldehydes/Propanone	
H.C. RAI, PRIYANKA, ANNAPURNA SHARMA, MEERA SHARMA and SARITA KUMARI	2267
Classical Approaches of Genetic Algorithms and Their Suitability	
LORENTZ JÄNTSCHI, SORANA D. BOLBOACA and RADU E. SESTRAS	2275
Oxidation of Oximes of Heterocyclic Ketones by Pyridinium Chlorochromate: A Kinetic and Mechanistic Study	
T.S. JANI BAI and S. SANTHI	2285
Kinetics and Mechanism of Pyridinium Chlorochromate Oxidation of Some N-Methyl-2,6-diphenylpiperidin-4-one Oximes	
T.S. JANI BAI and S. SANTHI	2290
Synthesis and Antimicrobial Study of 3,5,6-Trichloropyridine-2-yl(arylthio)acetate Derivatives	
L.S. SHET and A.R. SHELAR	2295
Determination of Diosmin by Visible Spectrophotometry	
P. UMA DEVI, K.V.V.V. SATYANARAYANA and C.S.P. SASTRY	2300
Determination of Domperidone and Omeprazole in Capsule Dosage form by UV/Visible Spectrophotometer	
V.P. PANDEY, S. VANJINATHAN and STELLA SINGH	2305
Influence of Traditional Processing on Some Compounds of Rose Hip ( <i>Rosa canina</i> L.) Fruits Collected From Habitat in Bursa, Turkey	
CIHAT TURKBEN, VILDAN UYLASER and BIGE INCEDAYI	2309
Adsorption Behaviour of Cr(III) Ions on Sepiolite	
Y. ASÇI and P. BERKAN	2319
Adsorption-Desorption Studies on Cr(VI) Using Non-Living Fungal Biomass	
MOUSUMI SEN and MANISHA GHOSH DASTIDAR	2331
Comparison of Olive Oils Derived from Certified Organic and Conventional Agricultural Methods	
OGUZ DOLGUN, GULCAN OZKAN and BUKET ERBAY	2339

Extraction of Adenosine from <i>Ganoderma lucidum</i> Using Novel Extraction Technologies N. YOSWATHANA, M.N. ESHTIAGHI and A.G. EBADI	2349
Measurement of Chemical and Non-Chemical Parameters of Three Native Pistachio Cultivars of Damghan Region (Iran) for Studying Spring Frost HOSSEIN AFSHARI, HOSSEIN HOKMABADI, ABDOLGHAPHAR EBADI and GHANBAR LAEE	2356
Determination of 107 Pesticide Residues in Citrus Fruits by Gas Chromatography/ Mass Spectrometry O. TAGA and B. BILGIN	2367
Effect of Streptozotocin on Biochemical Parameters in Rats CENNET RAGBETLI and EBUBEKIR CEYLAN	2375
Spectral Studies on Hydridophosphine Complexes of Pt-Group Metals: Part-III R.N. PANDEY, GUNJAN KUMARI and RAJNISH KUMAR SINGH	2379
Effect of 6-Benzylaminopurine, 2,4-Dichlorophenoxyacetic Acid and Indole-3-Butyric Acid on Micropropagation Stages of <i>Achillea biebersteinii</i> DANIAL KAHRIZI and MEHDI KAKAEI	2383
Evaluation of Nitrogen Fertilizer and Plant Density on Chemical Analysis of Corn Forage and Weed Population in an Agroforestry System S. GHANBARZADE, M.R. CHAICHI, S.M.B. HOSSEINI and A. SADEGHPOUR	2387
Chemical Analysis of Barley Under Different Irrigation and Fertilizing Systems S. MALEKI FARAHANI, M.R. CHAICHI*, D. MAZAHARI, R. TAVAKKOL AFSHARI and G. SAVAGHEBI	2397
Chemical Composition and Cytotoxic Activity of <i>Eruca sativa</i> L. Seeds Cultivated in Egypt NAGLAA M. NAZIF, AMIRA A.E. HABIB, WAFAA A.M. TAWFIK and RASMIA A. HASSAN	2407
Use of Gravimetric Separators for the Beneficiation of Gunduzler Chromite Ore U. MALAYOGLU, S. SEN and V.T. ENGIN	2417
Investigation of Nutrients Controlling Bacterial Regrowth in a PEX and Stainless Steel Plumbing Water System SANGCHUL NAM and YOON-JIN LEE	2427

## NOTES

Formulation and Characterization of Size Reduced Oral Extended Release Metformin Tablets OM PRAKASH TIWARI, SURYA PRAKASH B.N. GUPTA, NEERAJ UPMANYU, NAGESH CHIKMATH and V.S. DATTA MURTHY	2442
Synthesis of New Piperazino Oxadiazoles as Antiparkinsonian Agents RISHU AGARWAL, ALOK SAHAY and SUDHANSHU DWIVEDI	2445
UV Spectrophotometric Estimation of Phenytoin Sodium in Pure and Pharmaceutical Formulations MANGESH A. MANTRI and S. SHANMUKHAPPA	2447
Effect of <i>Leucas aspera</i> on Lead Acetate Induced Nephrotoxicity in Rats K. SOMA SEK HAR REDDY, Y. PADMANABHA REDDY, J. RAVINDRA REDDY, B. RAMESH, M. SRIRAMA CHANDRA, B. PRADEEP KUMAR and P. SANDHYA RANI	2450

Mixed-Ligand Complexes of $\text{Cu}^{2+}$ , $\text{Ni}^{2+}$ , $\text{Co}^{2+}$ , $\text{Zn}^{2+}$ with 2,2'-Bipyridine as a Primary Ligand and DL-2-Aminobutanedioic Acid as Secondary Ligand SHARDA SONI, MANU SIKARWAR and AVNISH SHARMA	2453
Undecyl 2,3-Dimethoxycinnamate from <i>Daphne oleoides</i> N. ULLAH	2456
Evaluation of Antioxidant Activity of Aerial Parts of <i>Hygrophila difformis</i> D.K. PAL, K. SAMANTA and P. MAITY	2459
Fatty Acid Composition of Some New Varieties of Oil Seeds SATISH INGALE and S.K. SHRIVASTAVA	2462
Investigation on Transition Metal Complexes with Sulphur and Nitrogen Containing Ligand Derived from Diethylamine MOHAMMAD TARIQUE and MOHAMMAD ASLAM	2465
Phytochemical Screening and Analgesic Activity of <i>Syzygium alternifolium</i> (Wild) V.V.S.S. APPALA RAJU, S. GANAPATY, S. RAO PIPPALLA and V. ASHA JYOTHI	2468
Phytochemical Studies on <i>Cissus Glauca</i> G.T. ABRAHAM and P.M. SHAFI	2471
Determination of Fatty Acids in Plant Seeds of Leguminosae Family from Arid Zone of Rajasthan ARUN ARORA and RAJENDRA SEN	2474
Conductometric Equilibrium Study of Methylmethionine Sulfonium Chloride J.J. VORA and L.S. BHUTADIYA	2477
Evaluation and Study of Minerals from <i>Leea macrophylla</i> Roxb (Leeaceae) K.D. JADHAO and M.P. WADEKAR	2480
Comparative Study of Vitamins Nutrient from <i>Marsilea quadrifolia</i> Linn. K.D. JADHAO and M.P. WADEKAR	2483
<b>CONTRIBUTORS TO THIS ISSUE</b>	ix



## Classical Approaches of Genetic Algorithms and Their Suitability

LORENTZ JÄNTSCHI†‡, SORANA D. BOLBOACA\*‡ and RADU E. SESTRAS‡

*"Iuliu Hatieganu" University of Medicine and Pharmacy Cluj-Napoca,  
13 E. Isac, 400023 Cluj-Napoca, Romania*

*Fax: (4)(264)593847; Tel: (4)(264)431697; E-mail: sbolboaca@umfcluj.ro*

Genetic algorithms derived from observations of nature and simulation of artificial selection of organisms with multiple loci that control a measurable trait. To date, genetic algorithms evolved into complex and strong informatics tools able to deal with hard problems of decision, classification, optimization and simulation. A series of studies reported biotechnology hard problems solved using genetic algorithms. In this context, the aim of the present article is to introduce genetic algorithms and to present their suitability for biotechnology hard problems. Important results are reported in the available literature that deal with the application of genetic algorithms for biotechnology process modelling.

**Key Words:** Genetic algorithms, Hard problems, Processes modelling, Kinetic models.

### INTRODUCTION

Any problem which has its complexity and the complexity of algorithms applied to find the optimum solution differ in terms of time (time complexity *e.g.*, the number of transitions from start to the end, hopefully with the correct answer) and space (space complexity *e.g.*, amount of random access memory required to the program for run) from one approach to another. A hard problem is one for which all algorithms that solve it are of high complexity. The problems with exponential complexity are also considered hard because even the best algorithm is used, it will probably be unusable on real-world instances<sup>1</sup>. If a problem is hard, then the search for the optimum solution often goes into out-of-time for real world applications. Besides, a large set of problems encountered in practice do not necessarily call for the optimum. Because most of the hard problems subsist from many years, for some of them one or several heuristics have already been formulated. These are rules of thumb recipes for solving a particular problem, usually based on common sense and avoiding obvious mistakes. Three heuristics applicable to a wide range of hard problems, known as meta-heuristics, were developed. All three are stochastic in nature and two of them are based on natural processes that have been taking place such as tabu search<sup>2</sup>, simulated annealing<sup>3</sup> and genetic algorithms<sup>4</sup>.

†Technical University of Cluj-Napoca, 103-105 Muncii Bvd, 400641 Cluj-Napoca, Romania.

‡University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, 3-5 Manastur, 400372 Cluj-Napoca, Romania.

In regards of genetic algorithms (GAs) history, the computer simulations of evolution started with the work of Nils Aall Barricelli<sup>5</sup>. Shortly later, the Australian quantitative geneticist Alex Fraser published a series of papers on simulation of artificial selection of organisms with multiple loci controlling a measurable trait<sup>6</sup>. Fraser's simulations included all of the essential elements of modern genetic algorithms.

Genetic algorithms (GAs) are adaptive heuristic search algorithm developed to mimic some of the processes observed in natural evolution with the idea to use this power of evolution to solve optimization problems. Genetic algorithms are designed to simulate processes in natural systems necessary for evolution, specially those follow the principles first laid down by Charles Darwin of survival of the fittest. Since in nature, competition among individuals for scanty resources results in the fittest individuals dominating over the weaker ones. Genetic algorithms are implemented as computer simulations in which a population of abstract representations (called chromosomes or the genotype of the genome) of a candidate solutions (called individuals, creatures, or phenotypes) subject to an optimization problem which evolves toward better solutions. Genetic algorithms simulates the survival of the fittest among individuals over consecutive generation for solving the problem. Each generation consists of a population of character strings analogous to the DNA chromosomes. Each individual represents a point in a search space and a possible solution. The individuals in the population are then made to go through a process of evolution. Genetic algorithms is based on an analogy with the genetic structure and behaviour of chromosomes within a population of individuals.

The genetic algorithms were applied to the hard problems from many scientific fields since popular in the early 1970s<sup>7</sup>: computer science and engineering<sup>8-12</sup>, agriculture<sup>13,14</sup>, medicine and chemistry<sup>15-19</sup>.

The aim of this study was to illustrate how genetic algorithm can be used to solve hard problems from the biotechnology field. The study includes a classical approach, which is minted to prepare the arena for genetic algorithms. Since is no unique approach to solve a problem, the comparison between different approaches were revealed.

## EXPERIMENTAL

Natural as well as controlled processes evolve through a mechanism. If it refers to chemistry, then the mechanism is about explaining the pathway of a reaction, whilst in biology is about explaining how a feature is created. Starting from a biochemistry approach<sup>20</sup>, the fast reaction between copper and thiosulfate ions was investigated<sup>21</sup>. The formulation of the kinetic problem is as follows: (i) two reactants (A and B) through forming of an intermediary X lead to the products (P), (ii) the intermediary X concentration can be correlated with an observed absorption of light intensity at a given wavelength (430 nm) using the well known Lambert-Beer relationship<sup>22</sup>, (iii) as it is in any general kinetics study, of particular interest is to obtain the reaction rates and partial orders.

Model formulation starts assigning of unknown reaction rates and orders for every elementary reaction of the process and writing of the reaction kinetics. Thus, for a reaction with pre-equilibrium, following chemical reactions and mathematical equations applied.;  $A + B \rightarrow X$  ( $d[X]/dt = k_0 \cdot [A]^{y_0} [B]^{y_1}$ );  $X \rightarrow A + B$  ( $d[X]/dt = -k_1 \cdot [X]^{y_2}$ );  $X \rightarrow P$  ( $d[X]/dt = -k_2 \cdot [X]^{y_3}$ ), where  $[\cdot]$  denote concentration in the aqueous solution. A mass conservation principle is applied if the process evolve in a controlled environment and starts from two initial concentration of reactants A and B, without addition of the reactants during the reaction. The following relationships between two instantaneous concentrations  $[\cdot]_1$  and  $[\cdot]_2$  are observed:  $[A]_1 - [A]_2 = [B]_1 - [B]_2 = [P]_2 - [P]_1 + [X]_2 - [X]_1$ .

The kinetic model (KM) can be state as: (i) chemical reaction:  $A+B \xrightleftharpoons[k_0]{k_1} X \xrightarrow{k_2} P$ ; (ii) a fundamental assumption:  $A = A(t)$ ,  $B = B(t)$ ,  $X = X(t)$ ,  $P = P(t)$  and mathematical equations (using A in place of  $[A]$  and so on):  $dA/dt = -k_0 A^{y_0} B^{y_1} + k_1 X^{y_2}$ ,  $dB/dt = -k_0 A^{y_0} B^{y_1} + k_1 X^{y_2}$ ,  $dX/dt = k_0 A^{y_0} B^{y_1} - k_1 X^{y_2} - k_2 X^{y_3}$  and  $E \sim X \iff \hat{E} = aX+b$  (from Lambert-Beer); (iii) mathematical inequalities:  $A, B, X, P \geq 0$  (concentrations);  $y_0, y_1, y_2, y_3 \geq 0$  (partial orders);  $k_0, k_1, k_2 \geq 0$  (reaction rates); (iv) experimental data:  $E = E(t)$ , experimental extinctions; (v) another fundamental assumption: during the reaction, the values  $k_0, k_1, k_2, y_0, y_1, y_2, y_3$  remains constant; (vi) the objective function:  $(aX+b-E)^2 = \min$ .

The kinetic model problem is a hard optimization problem, having the mathematical equations and inequalities, an objective function to be minimized and nine parameters to be determined. Fig. 1 presents a classical approach of solving algorithm<sup>23</sup>.




Fig. 1. Classical approach of a solving algorithm [Ref. 23]

The explanation of the classical approach of solving algorithm from Fig. 1 is as follows: (a) The initial values of the constants are stored into an array of data (which is optimized iteratively). (b) The iterative module uses the  $k_0...k_2, y_0...y_3$  values to generate a table with  $3^7$  ( $3 = \{-1;0;1\}$ ) rows and 7 columns ( $7 = \{k_0, k_1, k_2, y_0, y_1, y_2, y_3\}$ ). (c) Every row of the table generated in iterative module is used into the estimative module in order to estimated the compounds' concentration variation during time (reactants A and B, intermediary X and product P). (d) The objective module uses the  $\{A, B, X, P\}$  estimations to obtain the coefficients of regression  $\{a, b\}$ . (e) The values  $\{v_0...v_6\}$  which provide the smallest value of the objective function  $\Sigma(aX+b-E)^2$  are saved into  $\{vob_0...vob_6\}$  array. (f) At the end of a complete iteration of the iterative module ( $3^7$  iterations) the  $\{vob_0...vob_6\}$  values replace the old  $k_0...k_2, y_0...y_3$  values. (g) The cycle *Iterative module*  $\rightarrow$  *Estimative module*  $\rightarrow$  *Objective module* are repeated until the estimation of the unknown parameters  $k_0...k_2, y_0...y_3$  produce a stable approximation (after a iteration cycle their values does not vary significant).

**Method: The genetic algorithms approach:** In terms of a genetic algorithm, the kinetic model problem has: (a) The genetic code of a solution composed from seven genes  $\{k_0, k_1, k_2, y_0, y_1, y_2, y_3\}$  (Fig. 2). (b) An initial population that may represent a chosen number of individuals (let be for example of 100) randomly selected. (c) Selection process may give the chance to survival of the fittest for a given per cent of the individuals (let's say 50 %), allowing them to pass to the next generation based on the goodness of each individual that depends on its fitness, assessed by the objective function, which simultaneously obtain the 'a' and 'b' values (of remaining two parameters, Fig. 3). (d) Crossover. Individuals (two for example) are chose from the population using the selection operator. A double crossover along the bit strings is chose (usually randomly) then the values of the genes are exchanged up to this point and the two new offspring created from this mating are put into the next generation of the population. If there are recombine portions of good individuals, this process is likely to create even better individuals (Fig. 4). (e) Mutation. Introduces random modifications; with some low probability, a portion of the genes of the new individuals will have their values flipped, with the purpose of maintaining diversity within the population and to inhibit premature convergence (Fig. 5).




Fig. 2. Genetic code for a solution of kinetic model problem




Fig. 3. Natural selection for kinetic model problem: genotype, phenotype and fitness




Fig. 4. A double crossing over (involves the breakage and rejoining of parental chromosomes)




Fig. 5. Mutation of a genotype in kinetic model problem

## RESULTS AND DISCUSSION

**Biotechnology processes modeled using genetic algorithms:** As can be seen from the previous section, the genetic algorithm is easier to digest and implement and does not imply as many as classical algorithm do computations for iteration. A series of hard problems in biotechnology were solved using genetic algorithms, process kinetic modelling being just one of the genetic algorithm applications.

Lee *et al.*<sup>24</sup> reported parameter estimation using a hybrid of simplex and genetic algorithm by introducing the simplex method as an additional operator in the genetic algorithm. During the reproduction of each iteration step, the hybrid approach applies the simplex method to a top percentage of the population to produce new candidate solutions in the next generation. The remaining of the new population is generated using the genetic algorithm reproduction scheme (*i.e.*, selection, crossover and mutation). The genetic algorithm was applied for optimization of three kinetics reactions, when significant improvements of algebraic methods were obtained: (a) Carboxylation of phosphoenolpyruvate (PEP) to oxaloacetate (OAA) catalyzed by

PPC (P-enolpyruvate), when CO<sub>2</sub> is transformed to phosphate (Pi): CO<sub>2</sub> + PEP → OAA + Pi; (b) Transformation of adenosine tri-phosphate (ATP) to adenosine di-phosphate (ADP) in the presence of OAA transformed to PEP catalyzed by PCK (phosphoenolpyruvate carboxy-Kinase): OAA + ATP → PEP + ADP + CO<sub>2</sub>; (c) Transformation of phosphoenolpyruvate (PEP) to pyruvate (Pyr) in presence of ADP (transformed to ATP) catalyzed by pyruvate kinase (PyKi): PEP + ADP → Pyr + ATP.

Pizarro *et al.*<sup>25</sup> reported an evolution of the basic genetic algorithm adapted to the features of the model to explain the industrial fermentation growth rate of acetic fermentation. Each chromosome represents in their approach a possible combination of values of the five parameters to optimize, in binary code. There was imposed an allowed range of values for each parameter to adopt (as many values as allowed by the binary codification and the number of significant figures). The initial population was composed of randomly selected values for the parameters within the allowed ranges and codified into binary code. The evaluation program decodes the values of the parameters for each chromosome and then uses them to simulate a batch process with each sequence of parameters. The simulation algorithm solves a system of differential equations giving overall rates, viable biomass concentration and relationships between the product formation, substrate consumption and the cell growth using the Runge-Kutta algorithm. The initial concentrations were those of the representative sequence of the process and the initial viable biomass/total biomass ratio represents the parameters of the chromosome. There was an oxygen control in the simulation, because in the real process the oxygenation conditions are enough to satisfy the demanded amount of oxygen. The algorithm had two important stop conditions: when no real positive values for one concentration are obtained and when the process time in the simulation has reached the final process time of the representative sequence. A new generation with the same number of chromosomes is formed by applying reproduction, crossover and mutation operators. The chromosomes with the best fitting ability obtained the best value in the desirability function (*i.e.* closer to 1) and have more chances of being selected and copied into the next generation. Uniform crossover is used and the five best chromosomes of each generation pass unchanged to the next generation. These chromosomes are called elitist chromosomes. Twins and out-of-range chromosomes are disallowed by using a 'while' loop with filters. When some of these chromosomes are discovered after crossover, chromosomes also obtained by crossover substitute them and if they are discovered after mutation, they are replaced by the original chromosomes in the same positions but mutated again with the same chances of mutation. With this process, the mutability is not increased and the number of chromosomes remains constant. The process stops after five generations without changes higher than a fixed percentage of the mean response of the elitist chromosomes. The algorithm is completed five times each time when the program is run. A final run when the initial population is composed of the best chromosomes found in each of the previous

runs is performed. The acetic concentrations in the fermentators of the industrial plant of Vinagreras Riojanas SA (Logrono, Spain), obtained by NIR, were studied. The data were obtained for a period of 4 months without changes in the industrial parameters of the process, *i.e.* oxygenation conditions and temperature. The average temperature was 29.5 °C and the oxygenation conditions were enough to satisfy the oxygen demand and thus the oxygen became a non-limiting substrate. Nowadays, the fermentators of the industrial plant work discontinuously with charges. The batch bioreactors studied were always fed with white wine of the same origin. The process time was about 30-31 h and 218 complete sequences were obtained. An average concentration sequence was calculated by analyzing the data. This sequence is representative of the process to be modelled. The variability in the concentrations among the sequences is due to analytical errors and to factors that cannot be controlled in an industrial process (*i.e.* differences in the ethanol concentration of the wine among batch processes). Therefore, the model obtained with this sequence does not model this variance.

Guangzhu *et al.*<sup>26</sup> improved a simple genetic algorithm developing a hybrid genetic algorithm, which was used to estimate the kinetic parameters of polyesterification between dimer fatty acid and ethylene glycol. The work proved that the model developed by authors is useful for the polyesterification of dimer acid and ethylene glycol catalyzed by *p*-toluene sulfonic acid. The authors used 28.1 g (0.05 mol) of dimer fatty acid, 3.11 g (0.05 mol) of ethylene glycol and 0.5 % of *p*-toluene sulfonic acid as the catalyst. All were placed into a round bottom flask (three necks), which was equipped with a dephlegmator and a pipe for the nitrogen. Nitrogen was introduced into the flask to remove the oxygen and to prevent the oxidation of the materials. The flask was placed into an oil bath at 170 °C. After 0.5 h reaction, the nitrogen was stopped and vacuum pumping was used to remove the water from reactant. The reaction continued 8-10 h in vacuum. The acid value of the reactant was measured at certain reaction times during the progress of the reaction. The estimation of the parameters was carried out in three steps. First, the order of reaction was confirmed using the assumption of equal activity. Second, experiments were designed to estimate the parameters of rate constant of the reaction between carboxylic group on the monomer and hydroxyl group of the polymers. Excess monomer was supplemented into the reactant after it had reacted for several hours with the materials proportion of 1:1 and the reactions could be ignored except for the added monomer and the polymers. Finally, obtained values were introduced into the rate equations to obtain the values of reaction rates between carboxylic end of the monomer and hydroxyl group of the monomer and carboxyl on the polymers and hydroxyl on the polymers.

Moscovitch *et al.*<sup>27</sup> used the genetic algorithm in kinetic analysis of multiple proton transfer reactions. They demonstrated that the search of the rate constants can be fully automated using the genetic algorithm approach leading to a detailed kinetic analysis due to ability of identification of multiequilibria systems. A system

with inherent complexity made of seven independent variables consisting of a proton emitter (a pyranine molecule that eject a proton when excited by a photon), indicator (fluorescein, a pH indicator with two proton-binding sites represented by the oxyanion of the xanthene ring and the carboxylate of the benzene) and bicarbonate anion ( $\text{HCO}_3^-$ , a buffer molecule with unknown concentration that reacts with the proton but not generate a measurable signal) was investigated. The complexity of the studied system is comparable to the studying the protonation dynamics of an indicator attached to a protein<sup>28-31</sup>. The authors used a 100 mM NaCl aqueous solution supplemented with pyranine (20 mM) and fluorescein (10 mM), equilibrated with the air at two pH values (6.8 and 7.3) and subjected to a train of laser pulses (1-1.5 mJ/pulse; 10 Hz, 355 nm, 3 ns full-width half maximum). The absorption transients were recorded at 458 and 496 nm, where pyranine and fluorescein are, respectively, absorbing (the time resolutions were either 30 ns or 300 ns per data point and were converted into concentration units using the extinction coefficients  $24000 \text{ M}^{-1} \text{ cm}^{-2}$  for pyranine and  $50000 \text{ M}^{-1} \text{ cm}^{-2}$  for fluorescein). Two reactants of the system are directly observable. The rate constants of protonation of these reactants were measured (manual analysis) and the complexity of the system was increased by taking into consideration the concentration of the bicarbonate. The parameter's space was used in order to search for the uniqueness. The values of adjustable parameters were used to reconstruct the signal and to calculate the fittest. The first generation consisted of 100 phenotypes (randomly selected values for the adjustable parameters). The best-fit phenotype was cloned and replaces the worst-fitting one. The genetic algorithm was searching for the minimum of the fitness function in a seven-dimensional space. Each runs last 3000 generations from 2 to 6 h depending on the processor of the computer used. A stable solution in terms of no tendency to drift into a new set of adjustable parameters was search. The genetic algorithm was considered unique when the target signals were noiseless (the values derived by the program were identical to those used to create the signal). The results revealed that genetic algorithm is a reliable method for searching a solution of kinetics equations when the rate constants must be determines proving that the system work even if the concentration of one reactant is unknown. The utility of the genetic algorithm in solving chemical kinetic problems have also been proved by several researchers<sup>32-35</sup>.

Popelier *et al.*<sup>36</sup> modelled the mutagenic activity of 23 triazines and 24 halogenated hydroxyfuranones in order to generate significantly statistic valid quantitative models and to identify the active centre of the investigated compounds. A genetic algorithm was used to optimize the number of descriptors of the best model expressed as best possible coefficient of determination and leave-one-out cross validation coefficient. The genetic algorithm introduced a population of 256 randomly selected models and the cross-validation error was the fitness function at a mutation rate of 0.003 and for a maximum number of generations equal to 200. The analysis suggested a preferred mechanistic pathway for the initial hydroxylation of the triazines and elucidates the mechanistic ambivalence of hydroxyfuranones. A similar approach



was applied in investigation of steroid binding affinity and antibacterial activity of nitrofurans derivatives<sup>37</sup>. The genetic algorithm was used in order to select variable in the best performing model. Similar approach was used by Matsuda *et al.*<sup>38</sup> when the authors were able through their design system cyclopaedically to generate ion liquids structures corresponding to particular physical properties.

### Conclusion

Having a population of abstract representations (the genotype of the genome) of candidate solutions (phenotypes) genetic algorithm optimization problem evolves toward better solutions simulating the survival of the fittest among individuals over consecutive generation for solving the problem as living organisms evolve in nature.

The usage of genetic algorithm may reduce the algorithm complexity as was shown in kinetic model hard optimization problem.

Key hard problems solved using genetic algorithms include modelling industrial fermentation growth rate of acetic fermentation, heavy oil thermal cracking 3-lumping, fluid catalytic cracking unit main fractionator, reverse engineering of molecular mechanical machines, chemical kinetic problems and selection of descriptors used in quantitative structure-property/activity relationship models.

### ACKNOWLEDGEMENT

This work was in part supported by grants (ID-206/2007, ID-202/1.10.2007) of the National University Research Council, Romania.

### REFERENCES

1. E. Falkenauer, *Genetic Algorithms and Grouping Problems*, Wiley, New York (1998).
2. F. Glover, *Decision Sci.*, **8**, 156 (1977).
3. L. Davis, *Genetic Algorithms and Simulated Annealing*, San Francisco: M. Kaufmann (1987).
4. J. Bosworth, F. Norman, B.P. Zeigler, *Comparison of Genetic Algorithms with Conjugate Gradient Methods*, NASA Contractor Reports, CR-2093 (1972).
5. N.A. Barricelli, *Methodos*, 45 (1954).
6. A. Fraser, *Aust. J. Biol. Sci.*, **10**, 484 (1957).
7. J.H. Holland, In *Proceedings of the 1970 IEEE Symposium on Adaptive Processes Decision and Control: XVII*, ACM PRESS, New York (1970).
8. A.M. Barros, H.S. Lopes and A.L. Stelle, In *Proceedings of the 2007 IEEE Symposium on Computational Intelligence in Image and Signal Processing, CIISP 2007*, pp. 151-156 (2007).
9. J. Northern III and M. Ribeiro, In *Proceeding of the 2007 IEEE Region 5 Technical Conference, TPS*, Art. no. 4380385, pp. 223-227 (2007).
10. M.E. Aydemir, T. Gunel, S. Kargin, I. Erer and S. Kurnaz, In *Proceedings of 2nd International Conference on Recent Advances in Space Technologies*, Art. no. 1512654, pp. 684-688 (2005).
11. M. Sato, S. Matsumoto, Y. Teramoto and N. Adachi, *Trans. Japan. Soc. Artif. Intel.*, **16**, 324 (2001).
12. R. Neruda, *Neural Network World*, **11**, 267 (2001).
13. H. Muhlenbein and D. Schlierkamp-Voosen, *Evol. Comput.*, **1**, 335 (1993).
14. R. Salomon, *BioSystems*, **39**, 263 (1996).
15. P. Willett, *Trends Biotechnol.*, **13**, 516 (1995).
16. G. Jones, P. Willett, R.C. Glen, A.R. Leach and R. Taylor, *J. Mol. Biol.*, **267**, 727 (1997).

17. S. Kikuchi, D. Tominaga, M. Arita, K. Takahashi and M. Tomita, *Bioinformatics*, **19**, 643 (2003).
18. R. Leardi, *J. Chemometr.*, **15**, 559 (2001).
19. O. Roeva, T. Pencheva, S. Tzonkov, M. Arndt, B. Hitzmann, S. Kleist, G. Miksch, K. Friehs and E. Flaschel, *E. J. Biotechnol.*, **10**, 593 (2007).
20. L. Jäntschi, *Leonardo J. Sci.*, **2**, 1 (2003).
21. C.E. Stoenoiu, S.D. Bolboaca and L. Jäntschi, In Proceeding of the 4th ICAMC, Plovdiv, Bulgaria, p. 511 (2007).
22. P.W. Atkins, *Physical Chemistry*, Oxford: Oxford University Press, edn. 5 (1994).
23. L. Jäntschi and S.D. Bolboaca, Processes at Phases Interface: Mathematical Modelling, Numerical Optimization, Web Implementation, with Applications for Separation and Characterization of Biological Active Compounds Series (MEC/5892/18.09.2006)-Research Synthesis for 2008 year (in Romanian), UEFISCSU-ET108 (2008).
24. B. Lee, J. Yen, L. Yang and J.C. Liao, *Biotechnol. Bioeng.*, **62**, 722 (1999).
25. C. Pizarro, J.M. Gonzalez-Saiz and D. Garrido-Vidal, *J. Chemometr.*, **17**, 453 (2003).
26. F. Guangzhu, L. Fasong, L. Heping, Q. Hai and C. Yingde, *Chem. Eng. Technol.*, **29**, 740 (2006).
27. D. Moscovitch, O. Noivirt, A. Mezer, E. Nachliel, T. Mark, M. Gutman and G. Fibich, *Biophys. J.*, **87**, 47 (2004).
28. E. Nachliel, M. Gutman, S. Kiryati and N.A. Dencher, *Proc. Natl. Acad. Sci. (USA)*, **93**, 10747 (1996).
29. E. Nachliel and M. Gutman, *Biochim. Biophys. Acta*, **1514**, 33 (2001).
30. E. Shimoni, E. Nachliel and M. Gutman, *Biophys. J.*, **64**, 480 (1993).
31. R. Yam, E. Nachliel and M. Gutman, *J. Am. Chem. Soc.*, **110**, 2636 (1988).
32. B. Filipic, I. Zun and M. Perpar, *Int. J. Hum. Comput. Stud.*, **53**, 517 (2000).
33. C. Hongqing, Y. Jingxian, K. Lishan, C. Yuping and C. Yongyan, *Comput. Chem.*, **23**, 143 (1999).
34. D.B. Terry and M. Messina, *J. Chem. Inf. Comput. Sci.*, **38**, 1232 (1998).
35. C. Viappiani, G. Bonetti, M. Carelli, F. Ferrati and A. Sternieni, *Rev. Sci. Instrum.*, **69**, 270 (1998).
36. P.L.A. Popelier, P.J. Smith and U.A. Chaudry, *J. Comput.-Aided. Mol. Des.*, **18**, 709 (2004).
37. P.J. Smith and P.L.A. Popelier, *J. Comput.-Aided. Mol. Des.*, **18**, 135 (2004).
38. H. Matsuda, H. Yamamoto, K. Kurihara and K. Tochigi, *Fluid Phase Equilib.*, **261**, 434 (2007).

(Received: 30 June 2009;

Accepted: 1 December 2009)

AJC-8119

**CONTRIBUTORS TO THIS ISSUE**

(Chief Authors' name are in bold)

- Abd Al-Rahman, Abd Al-Moujod M., 2179  
**Abdalrazaq, Eid A., 2179**  
 Abraham, G.T., 2471  
**Adak, M.R., 2099**  
**Afshari, Hossein, 2356**  
 Afshari, R. Tavakkol, 2397  
 Agarwal, Kavita, 1822  
**Agarwal, Rishu, 2445**  
 Agarwal, S., 2051  
 Akhavan, Reza, 1925  
 Ali, Qasim, 1779  
**Alulu, S., 2129**  
 Amutha, P. Sahaya, 1918  
**Ankara, Huseyin, 2073**  
**Anwar, Farooq, 1779**  
**Appala Raju, V.V.S.S., 2468**  
 Arici, M., 1879  
**Arora, Arun, 2474**  
 Arshadi, S., 1907  
**Arslantas, Ali, 1709**  
**Asçi, Y., 2319**  
 Ashfaq, M., 1787  
**Ashnagar, A., 2058**  
 Ashraf, Muhammad, 1779  
 Aslam, Aftab, 1957  
**Aslam, Mohammad, 2031, 2465**  
 Assefipour, Reza, 2214, 2221, 2225  
**Atashi, Hossein, 1925**  
 Atsuta, Yoichi, 1693  
**Avci, Halim, 1829**  
 Azimi, Ali Akbar, 1665
- Babu, Ch. Suresh, 1822  
 Badeli, A., 1907  
 Bag, Swarnendu, 2147  
 Bano, Shaista, 1957  
 Barik, B.B., 1896  
 Basaran, Davut, 2081  
**Bekhradnia, A.R., 1907**  
 Berkan, P., 2319  
 Bhanja, S.B., 1896  
 Bhaumik, U., 2051  
**Bhowmik, Manas, 2147**  
**Bhutadiya, L.S., 2477**  
**Bi, Cai Feng, 1689**
- Bilgin, B., 2367**  
 Bilici, Adil Çagdas, 1853  
**Bolboaca, Sorana D., 2275**  
 Bose, A., 2051  
 Bozdemir, Yurdanur, 2013  
 Bruce, J.M., 2058  
 Buttrus, Nabeel H., 2179
- Cermak, R., 1982  
 Çetin, Mehmet, 1845  
 Çetin, Murat, 1845  
 Ceylan, Ebubekir, 2375  
**Chaichi, M.R., 2387, 2397**  
 Chakrapani, Ch., 1822  
 Champaneri, V.A., 1799  
 Chandanadevi, Bhupatiraju A.S., 1684  
 Chandra, A., 1675  
 Chandra, M. Srirama, 2450  
 Chang, Jun, 2018, 2172  
 Chang, Qiaowen, 1935  
 Chang, Zhi-Xian, 2207  
 Chatterjee, B., 2051  
 Chattopadhyay, Dipankar, 2147  
 Chen, M.L., 2231  
 Chen, X.H., 1889  
 Chen, Xizhu, 1935  
 Cheng, Zhao-Li, 1973, 1977  
 Chikmath, Nagesh, 2442  
**Chowdary, K.P.R., 1723**  
 Contractor, K.S., 2244
- Dai, Yun, 1705  
**Daimon, Hiroyuki, 1693**  
**Darafsheh, Mohammad Reza, 1966**  
 Das, A., 2051  
 Das, Sanchita, 2147  
 Dastidar, Manisha Ghosh, 2331  
 Demirci, M., 1879  
 Desai, S.A., 2244  
 Deshmukh, J.H., 1760  
**Deshpande, M.N., 1760**  
 Devanna, N., 1835  
 Dolgun, Oguz, 2339  
 Dwivedi, Sudhanshu, 2445

- Ebadi, A.G., 2349  
 Ebadi, Abdolghaphar, 2356  
 Ebrahimzadeh, M.A., 1907  
 Elsaid, A.G. Fathi, 1736  
 Emami, J., 1907  
 Engin, V.T., 2417  
 Erbay, Buket, 2339  
**Ergün, Yavuz, 1853**  
 Eshtiaghi, M.N., 2349  
 Eskandari, Habibollah, 2106
- Faisal, Muhammad, 1693  
 Fan, Peng, 1705  
**Fan, Yuhua, 1689**  
 Farahani, S. Maleki, 2397  
 Farhadi, K., 2165  
 Feiziasl, V., 1700  
 Feng, Shaoping, 2018, 2172
- Ganapaty, S., 2468  
**Gencoglu, Efe N., 1944**  
 Ghanbarzade, S., 2387  
 Gharacheh, Mohsen Ahmadi, 1949  
 Ghosh, A., 2051  
 Ghosh, Lakshmi K., 2147  
 Ghosh, Samir, 1835  
 Göçmentürk, Mustafa, 1853  
**Gouranlou, Farideh, 1873**  
 Gülle, Sibel, 1853  
**Gumus, T., 1879**  
**Gupta, O.D., 2035, 2043**  
**Gupta, Surya Prakash B.N., 2442**
- Habib, Amira A.E., 2407  
 Haghghi, Mohammad, 2106  
**Hassan, Hamed H.A.M., 2117**  
 Hassan, Rasmia A., 2407  
 Hokmabadi, Hossein, 2356  
 Hosseini, S.M.B., 2387  
 Hosseinzadeh, Marjan, 1949  
 Hrcirik, J., 1982  
**Hu, Qiu-Fen, 1705**  
**Huang, Zhangjie, 2018, 2172**  
 Hussain, Abdullah Ijaz, 1779
- Imran, Muhammad, 1993**  
 Incedayi, Bige, 2309  
**Ingale, Satish, 2462**
- Iqbal, Javed, 1993  
 Iqbal, Nida, 1993  
**Irez, Gazi, 1813**  
 Ismail, Mahmoud Hasan, 2187
- Jaafarzadeh, Nematollah, 1665, 1751  
 Jadhao, K.D., 2480, 2483  
 Janacova, D., 1982  
 Jani Bai, T.S., 2285, 2290  
 Jäntschi, Lorentz, 2275  
**Jin, Tong-Shou, 1973, 1977**  
 Jyothi, V. Asha, 2468
- Kahrizi, Danial, 2383**  
 Kakaei, Mehdi, 2383  
 Kale, Bhima, 1835  
**Kar, R.K., 1896**  
 Karadia, Chanchal, 2035  
 Karimi, S., 1700  
**Kaushik, R.D., 2090, 2136, 2155**  
 Kaya, Yunus, 1813  
**Keyvanfard, Mohsen, 1729**  
 Khan, Parwaz, 1957  
 Khan, S. Razi, 1787  
 Krishnaraju, Alluri V., 1684  
 Kumar, A. Sanjeev, 1835  
 Kumar, Ajay, 2090, 2155  
 Kumar, Ashok, 1661  
 Kumar, B. Pradeep, 2450  
 Kumar, Dharmendra, 2155  
 Kumar, Rajeev, 1769  
 Kumar, Tarun, 2136  
 Kumari, Anita, 2251  
 Kumari, Gunjan, 2379  
**Kumari, Kalpna, 1885**  
 Kumari, Reeta, 2251  
 Kumari, Sarita, 2251, 2267
- Lae, Ghanbar, 2356  
**Lee, Yoon-Jin, 2427**  
 Li, Chunying, 1935  
**Li, De-Liang, 2207**  
**Li, Gan-Peng, 1705**  
 Li, Tao, 1705  
 Li, Tong-Shuang, 1973, 1977  
 Liu, Li-Bin, 1977  
**Liu, Weiping, 1935**

- Mahmood, Zaid, 1993  
 Maity, P., 2459  
**Maity, T.K., 1841**  
 Malayoglu, U., 2417  
 Malik, Rajdeep, 2090, 2155  
 Mandil, H., 2129  
 Mandil, Hasna, 2187, 2195  
 Manila, 2090  
**Mantri, Mangesh A., 2447**  
 Marghmaleki, Mehran Asadi, 1939  
**Mathiyalagan, N., 1918**  
**Matter, A.B. Hamdy, 1736**  
 Mazaheri, D., 2397  
 Meena, 2043  
 Mehta, G.N., 1835  
 Messripour, Manochehr, 1939  
 Minaei, E., 2165  
**Modaresi, Mehrdad, 1939**  
 Moghaddas, Jafarsadegh, 2106  
 Moghani, Ali, 1966  
 Mohapatra, P., 1896  
 Mohapatra, S., 1896  
 Mohd, Ayaz, 1957  
**Mokrejs, P., 1982**  
**Mosavi, S.B., 1700**  
 Murthy, V.S. Datta, 2442  
 Mutlu, Hasene, 1813
- Naik, Satyanarayan, 1808**  
**Najafi, Hossein, 2214, 2221, 2225**  
 Nam, Sangchul, 2427  
**Nazif, Naglaa M., 2407**  
 Necefoglu, Hacali, 1709
- Oh, W.C., 2231**  
 Okay, Gürol, 1853  
 Oktav, Mehmet, 1944  
**Okumus, Veysi, 2081**  
 Onay, Ahmet, 2081  
 Oral, Elif, 2081  
**Ozkan, Gulcan, 2339**  
 Ozsunar, A., 1879  
**Öztürk, Serkan, 1845**
- Pahlevan, R., 1907  
**Pal, D.K., 2459**  
**Pal, T.K., 2051**  
**Pandey, R.N., 1657, 1661, 2379**
- Pandey, V.P., 2305**  
**Parmar, R.G., 1717**  
 Patel, R.N., 2244  
 Pathak, Prince, 1773  
**Pathak, Vishal, 1773**  
 Pazoki, Maryam, 1751  
 Pippalla, S. Rao, 2468  
 Prasad, Mahender, 1822  
 Priyanka, 2267
- Qajar, F., 1907  
**Qiu, G.Z., 1889**
- Ragbetli, Cennet, 2375**  
 Raghav, A.K., 1675  
**Raghuvanshi, Shiv Pratap, 1675**  
**Rai, H.C., 2251, 2267**  
 Rajae, M., 1907  
 Rajapandi, R., 1841  
**Rajeev, Rekha, 1769**  
**Ramadan, Abdul Aziz, 2187, 2195**  
 Ramanatham, J., 1835  
 Ramesh, B., 2450  
 Rani, P. Sandhya, 2450  
 Rao, Chirravuri V., 1684  
**Rao, G. Naga Bhushana, 2027**  
**Rao, K. Somasekhara, 1822**  
 Rao, Y. Ramachandra, 1808  
 Reddy, B.V. Rajeswara, 1822  
 Reddy, C. Umamaheswara, 1723  
 Reddy, J. Ravindra, 2450  
**Reddy, K. Soma Sekhar, 2450**  
 Reddy, Y. Padmanabha, 2450  
 Rizk, N.M.H., 1736  
 Rollas, Sevim, 1944  
 Rout, Prasant K., 1808
- Sabounchei, Seyyed Javad, 1949**  
**Sabzi, R.E., 2165**  
 Sadeghpour, A., 2387  
**Saeed, R., 1787**  
 Sahay, Alok, 2445  
 Sahebi-shahemabadi, Mehdi, 1925  
 Sakur, A.A., 2129  
 Samanta, K., 2459  
**Santhi, S., 2285, 2290**  
 Sastry, C.S.P., 2300  
 Satyanarayana, K.V.V.V., 2300

- Savaghebi, G., 2397  
 Savithri, Mantena G., 1684  
 Seenivasan, P., 1723  
**Sen, Mousumi, 2331**  
 Sen, Rajendra, 2474  
**Sen, S., 2417**  
 Sengupta, P., 2051  
 Sestras, Radu E., 2275  
 Shad, Y., 1907  
**Shafi, P.M., 2471**  
 Shah, S.S., 1717  
 Shanmukhappa, S., 2447  
 Sharma, Annapurna, 2251, 2267  
 Sharma, Avnish, 2453  
 Sharma, Meera, 2251, 2267  
**Shayesteh, Keivan, 2106**  
 Shelar, A.R., 2295  
 Shelat, C.D., 1745  
**Shet, L.S., 2295**  
 Shrivastava, S.K., 2027, 2462  
**Siddiqi, K.S., 1957**  
 Sikarwar, Manu, 2453  
**Singh, B.R., 1864**  
 Singh, D.P., 1661  
 Singh, Deep Sorabh, 1864  
 Singh, Jagadish, 1841  
 Singh, Jyoti, 2043  
 Singh, Monal, 1773  
 Singh, Prabha, 2136  
 Singh, Rajnish Kumar, 2379  
 Singh, Renu, 1675  
 Singh, Stella, 2305  
 Sinha, Jaydeep, 2147  
**Soni, Sharda, 2453**  
**Soundararajan, R., 1835**  
 Sozmen, Mahmut, 1709  
 Sundriyal, Purnima, 2136  
 Taga, O., 2367  
**Takdastan, Afshin, 1665, 1751**  
 Tarique, Mohammad, 2031, 2465  
 Tawfik, Wafaa A.M., 2407  
 Tiwari, Om Prakash, 2442  
**Trimurtulu, Golakoti, 1684**  
 Turkben, Cihat, 2309  
**Ullah, N., 2456**  
**Uma Devi, P., 2300**  
 Upmanyu, Neeraj, 2442  
**Uylaser, Vildan, 2309**  
 Vanjinathan, S., 2305  
 Vasek, V., 1982  
**Vashi, R.T., 1745, 1799, 2244**  
 Venugopal, S., 1835  
 Vora, J.J., 2477  
**Wadekar, M.P., 2480, 2483**  
 Wang, Ai-Qing, 1973, 1977  
 Weiping, Liao, 1859  
**Xiong, Chunhua, 2003**  
 Yaman, Mehmet, 1829  
 Yang, Jun, 1689  
 Yang, Zhongqi, 2018, 2172  
 Yao, Caiping, 2003  
 Yazdani, O., 1907  
 Ye, Qingsong, 1935  
 Yigmatepe, Ergul, 1829  
**Yoswathana, N., 2349**  
 You, Tingting, 1689  
 Yu, Yao, 1935  
**Yucebilgic, Guzide, 2013**  
**Yukui, Rui, 1859**  
 Zhang, F.J., 2231  
 Zhang, Jun, 2172  
 Zhang, Ling, 2207  
 Zhang, Ni, 1935  
 Zhang, Xin, 1705  
 Zhang, Zhi-Jun, 2207  
 Zheng, Zhanwang, 2003  
 Zhou, H.B., 1889